



**Queensland University of Technology**  
Brisbane Australia

This may be the author's version of a work that was submitted/accepted for publication in the following source:

Inoue, Akihiko, Nakano, Yumiharu, & Anh, Vo  
(2007)  
Binary Market Models With Memory.  
*Statistics and Probability Letters*, 77(3), pp. 256-264.

This file was downloaded from: <https://eprints.qut.edu.au/21838/>

**© Consult author(s) regarding copyright matters**

This work is covered by copyright. Unless the document is being made available under a Creative Commons Licence, you must assume that re-use is limited to personal use and that permission from the copyright owner must be obtained for all other uses. If the document is available under a Creative Commons License (or other specified license) then refer to the Licence for details of permitted re-use. It is a condition of access that users recognise and abide by the legal requirements associated with these rights. If you believe that this work infringes copyright please provide details by email to [qut.copyright@qut.edu.au](mailto:qut.copyright@qut.edu.au)

**Notice:** *Please note that this document may not be the Version of Record (i.e. published version) of the work. Author manuscript versions (as Submitted for peer review or as Accepted for publication after peer review) can be identified by an absence of publisher branding and/or typeset appearance. If there is any doubt, please refer to the published source.*

<https://doi.org/10.1016/j.spl.2006.07.007>

# BINARY MARKET MODELS WITH MEMORY

AKIHIKO INOUE, YUMIHARU NAKANO AND VO ANH

ABSTRACT. We construct a binary market model with memory that approximates a continuous-time market model driven by a Gaussian process with stationary increments. We give a sufficient condition for the binary model to be arbitrage-free. In a case when arbitrage opportunities exist, we present the rate at which the arbitrage probability tends to zero.

## 1. INTRODUCTION

Let  $T \in (0, \infty)$ . We consider the stock price process  $(S_t)_{0 \leq t \leq T}$  that is governed by the stochastic differential equation

$$dS_t = S_t(bdt + \sigma dY_t), \quad 0 \leq t \leq T, \quad (1.1)$$

where  $\sigma$  and the initial value  $S_0$  are positive constants, and  $b \in \mathbf{R}$ . In the classical Black-Scholes model, Brownian motion is used as the driving noise process  $Y$ , and the resulting price process  $S$  becomes Markovian. In Anh and Inoue (2005), Anh et al. (2005) and Inoue et al. (2006), the Gaussian process  $Y_t = B_t - \int_0^t \left\{ \int_{-\infty}^s p e^{-(q+p)(s-u)} dB_u \right\} ds$ ,  $0 \leq t \leq T$ , which has stationary increments, is used instead as the driving noise process  $Y$  in (1.1), where  $p$  and  $q$  are real constants such that  $0 < q < \infty$ ,  $-q < p < \infty$ , and  $(B_t)_{t \in \mathbf{R}}$  is a one-dimensional Brownian motion defined on a probability space  $(\Omega, \mathcal{F}, P)$  satisfying  $B_0 = 0$ . The parameters  $p$  and  $q$  describe the memory of  $Y$ , and the resulting stock price process  $S$  becomes non-Markovian.

We write  $(\mathcal{F}_t)_{0 \leq t \leq T}$  for the  $P$ -augmentation of the filtration generated by the process  $(Y_t)_{0 \leq t \leq T}$ . The theory of innovation processes as described in Liptser and Shirayev (2001) tells us that  $Y$  is an  $(\mathcal{F})$ -semimartingale (cf. Anh and Inoue, 2005, Theorem 3.1) though the above representation itself is not a semimartingale representation of  $Y$  since  $(B_t)$  is not  $(\mathcal{F}_t)$ -adapted. In fact, using the prediction theory for  $Y$  which is developed in Anh et al. (2005), the following explicit semimartingale representation of  $Y$  is obtained in Inoue et al. (2006):

$$Y_t = W_t - \int_0^t \left\{ \int_0^s l(s, u) dW_u \right\} ds, \quad 0 \leq t \leq T, \quad (1.2)$$

where  $(W_t)_{0 \leq t \leq T}$  is a one-dimensional Brownian motion, called the *innovation process*, satisfying  $\sigma(W_s : 0 \leq s \leq t) = \sigma(Y_s : 0 \leq s \leq t)$  for  $0 \leq t \leq T$  and  $l(t, s)$  is a bounded Volterra kernel given explicitly by

$$l(t, s) = p e^{-(p+q)(t-s)} \left\{ 1 - \frac{2pq}{(2q+p)^2 e^{2qs} - p^2} \right\}, \quad 0 \leq s \leq t \leq T. \quad (1.3)$$

---

2000 *Mathematics Subject Classification* Primary 91B28, 60G10; Secondary 60F17.

*Date:* May 19, 2006.

*Key words and phrases.* Financial market with memory, binary market, arbitrage.

This work is partially supported by the Australian Research Council grant DP0345577.

Thus the process  $Y$  has the virtue that it simultaneously possesses the property of a stationary increments process and the simple semimartingale representation (1.2) with (1.3). The market model driven  $Y$  is arbitrage-free and complete since the process  $Y$  becomes a Brownian motion under a suitable probability measure (see Anh and Inoue, 2005, Section 3).

As is well known, binary approximation of the Black-Scholes model plays a very important role for the model in many ways. Sottinen (2001) constructed a binary market model that approximates the market model driven by fractional Brownian motion, and investigated the arbitrage opportunities in the binary model. In this paper, we construct a binary market model with memory that approximates the continuous-time market model driven by  $Y$  in (1.2). However, rather than considering the special kernel  $l(t, s)$  in (1.3), we take a general bounded measurable Volterra kernel  $l(t, s)$ . We remark that any centered Gaussian process  $Y = (Y_t)_{0 \leq t \leq T}$  that is equivalent to a Brownian motion has a canonical representation of the form (1.2) with  $l(t, s)$  satisfying square integrability (see Hida and Hitsuda, 1991, Chapter VI). Thus, in this paper, we consider a subclass consisting of  $Y$  for which  $l(t, s)$  is bounded. As in Sottinen (2001), the key feature to the construction of the approximating binary market is to prove a Donsker-type theorem for the process  $Y$  (Theorem 2.1).

As stated above, the market driven by  $Y$  in (1.2) with (1.3) is arbitrage-free unlike that driven by fractional Brownian motion. However, the approximating binary market model may admit arbitrage opportunities. We consider conditions for their existence or non-existence. We also study the rate at which the arbitrage probability tends to zero.

## 2. A DONSKER-TYPE THEOREM

Let  $T \in (0, \infty)$ . In what follows, we write  $C = C_T$  for positive constants, depending on  $T$ , which may not be necessarily equal to each other. Let  $n \in \mathbf{N}$ . In Sections 2 and 3, we write  $\sum_{s \leq t} X_s = \sum_{i=1}^{\lfloor nt \rfloor} X_{(i/n)}$ ,  $\prod_{s \leq t} X_s = \prod_{i=1}^{\lfloor nt \rfloor} X_{(i/n)}$ .

Let  $l(t, s)$  be a bounded measurable function on  $[0, T] \times [0, T]$  that vanishes whenever  $s > t$ . Let  $W = (W_t)_{0 \leq t \leq T}$  be a one-dimensional Brownian motion on a probability space  $(\Omega, \mathcal{F}, P)$ . We define the process  $Y = (Y_t)_{0 \leq t \leq T}$  by (1.2).

We put, for  $t, u \in [0, T]$ ,

$$z(t, u) := \int_u^t l(s, u) ds, \quad y(t, u) := 1 - z(t, u).$$

Then both  $z(t, u)$  and  $y(t, u)$  are bounded and continuous on  $[0, T] \times [0, T]$ , and it holds that  $Y_t = \int_0^t y(t, u) dW_u$  for  $0 \leq t \leq T$ . Let  $C$  be a positive constant satisfying, for  $(t_1, u), (t_2, u) \in [0, T] \times [0, T]$ ,

$$|z(t_1, u) - z(t_2, u)| = |y(t_1, u) - y(t_2, u)| \leq C|t_1 - t_2|. \quad (2.1)$$

Let  $\{\xi_i\}_{i=1}^\infty$  be a sequence of i.i.d. random variables with  $E[\xi_1] = 0$ ,  $E[(\xi_1)^2] = 1$  and  $E[(\xi_1)^4] < \infty$ . We define the process  $W^{(n)} = (W_t^{(n)})_{0 \leq t \leq T}$  by

$$W_t^{(n)} := \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor nt \rfloor} \xi_i, \quad 0 \leq t \leq T,$$

where  $\lfloor x \rfloor$  denotes the greatest integer not exceeding  $x$ . The process  $W^{(n)}$  converges weakly to  $W$  in the Skorohod space by Donsker's theorem (see, e.g., Billingsley,

1968, Theorem 16.1). We define the process  $Y^{(n)} = (Y_t^{(n)})_{0 \leq t \leq T}$  by

$$Y_t^{(n)} := \int_0^t y\left(\frac{\lfloor nt \rfloor}{n}, s\right) dW_s^{(n)}, \quad 0 \leq t \leq T.$$

Then it follows that  $Y_t^{(n)} = n^{-1/2} \sum_{i=1}^{\lfloor nt \rfloor} y(\lfloor nt \rfloor/n, i/n) \xi_i$  for  $0 \leq t \leq T$ .

Here is the Donsker-type theorem for  $Y$ .

**Theorem 2.1.** *The process  $Y^{(n)}$  converges weakly to  $Y$  as  $n \rightarrow \infty$ .*

*Proof.* We first show that the finite-dimensional distributions of  $Y^{(n)}$  converge to those of  $Y$  as  $n \rightarrow \infty$ . Thus, for  $a_1, \dots, a_d \in \mathbf{R}$  and  $t_1, \dots, t_d \in [0, T]$ , we show that  $X^{(n)}$  converges to a normal distribution with variance  $\text{Var}(X)$ , where  $X^{(n)} := \sum_{k=1}^d a_k Y_{t_k}^{(n)}$  and  $X := \sum_{k=1}^d a_k Y_{t_k}$ . We have

$$\begin{aligned} \text{Var}(X^{(n)}) &= \sum_{k,l=1}^d a_k a_l \frac{1}{n} \sum_{i=1}^{\lfloor n(t_k \wedge t_l) \rfloor} y\left(\frac{\lfloor nt_k \rfloor}{n}, \frac{i}{n}\right) y\left(\frac{\lfloor nt_l \rfloor}{n}, \frac{i}{n}\right) \\ &= \sum_{k,l=1}^d a_k a_l \int_0^{\frac{\lfloor n(t_k \wedge t_l) \rfloor}{n}} y\left(\frac{\lfloor nt_k \rfloor}{n}, \frac{\lfloor ns \rfloor + 1}{n}\right) y\left(\frac{\lfloor nt_l \rfloor}{n}, \frac{\lfloor ns \rfloor + 1}{n}\right) ds, \end{aligned}$$

where  $t \wedge s := \min(t, s)$ . The function  $(t_1, t_2, u) \mapsto y(t_1, u)y(t_2, u)$  is continuous, whence uniformly continuous, on the compact set  $[0, T]^3$ . From this and the fact that  $0 \leq t - (\lfloor nt \rfloor/n) < 1/n$ , we see that

$$\lim_{n \rightarrow \infty} \text{Var}(X^{(n)}) = \sum_{k,l=1}^d a_k a_l \int_0^{t_k \wedge t_l} y(t_k, s)y(t_l, s) ds = \text{Var}(X). \quad (2.2)$$

We may assume  $\text{Var}(X) > 0$ . For, otherwise, (2.2) implies that  $X^{(n)}$  converges to  $X = 0$  in law. We put  $b_i^{(n)} := \sum_{k=1}^d a_k y(\lfloor nt_k \rfloor/n, i/n)$  and  $X_i^{(n)} := n^{-1/2} b_i^{(n)} \xi_i$  for  $n, i = 1, 2, \dots$ . Then we have  $X^{(n)} = \sum_{i=1}^{\lfloor nT \rfloor} X_i^{(n)}$  for  $n = 1, 2, \dots$ . We need to show the following Lindeberg's condition: for every  $\epsilon > 0$ ,

$$\lim_{n \rightarrow \infty} \sum_{i=1}^{\lfloor nT \rfloor} E \left[ (X_i^{(n)})^2 \mathbf{1}_{\{|X_i^{(n)}| > \epsilon \sigma^{(n)}\}} \right] = 0, \quad (2.3)$$

where  $\sigma^{(n)} := \sqrt{\text{Var}(X^{(n)})}$ . Choose a positive constant  $M$  satisfying  $|b_i^{(n)}| \leq M$  for  $n, i = 1, 2, \dots$ . Then since  $|X_i^{(n)}| \leq M n^{-1/2} |\xi_i|$ , we have

$$\begin{aligned} \sum_{i=1}^{\lfloor nT \rfloor} E \left[ (X_i^{(n)})^2 \mathbf{1}_{\{|X_i^{(n)}| > \epsilon \sigma^{(n)}\}} \right] &\leq \sum_{i=1}^{\lfloor nT \rfloor} E \left[ (M n^{-1/2} \xi_i)^2 \mathbf{1}_{\{|M n^{-1/2} \xi_i| > \epsilon \sigma^{(n)}\}} \right] \\ &= \sum_{i=1}^{\lfloor nT \rfloor} M^2 n^{-1} E \left[ (\xi_i)^2 \mathbf{1}_{\{|\xi_i| \geq M^{-1} \sigma^{(n)} \sqrt{n}\}} \right] \leq M^2 T E \left[ (\xi_1)^2 \mathbf{1}_{\{|\xi_1| \geq M^{-1} \sigma^{(n)} \sqrt{n}\}} \right]. \end{aligned}$$

We obtain (2.3) from this. By (2.3) and (2.2), we can apply the central limit theorem (cf. Billingsley, 1968, Theorem 7.2), so that  $X^{(n)}$  converges to  $X$  in law, as desired.

Next we show that, for  $0 \leq t_1 \leq t \leq t_2 \leq T$  and  $n = 1, 2, \dots$ ,

$$E \left[ |Y_t^{(n)} - Y_{t_1}^{(n)}|^2 |Y_{t_2}^{(n)} - Y_t^{(n)}|^2 \right] \leq C |t_2 - t_1|^2. \quad (2.4)$$

The theorem follows from this and Theorem 15.6 of Billingsley (1968). However, if  $t_2 - t_1 < 1/n$ , then either  $t_1$  and  $t$  or  $t$  and  $t_2$  lie in the same subinterval  $[\frac{m}{n}, \frac{m+1}{n})$  for some  $m$ , whence the left hand side of (2.4) is zero. Therefore we may assume that  $t_2 - t_1 \geq 1/n$ .

We show that

$$E \left[ |Y_t^{(n)} - Y_s^{(n)}|^4 \right] \leq C|t - s|^2 \quad (2.5)$$

for  $t, s$  and  $n$  satisfying

$$0 \leq s < t \leq T, \quad t - s \geq 1/n. \quad (2.6)$$

This implies (2.4) under the condition  $t_2 - t_1 \geq 1/n$  since

$$\begin{aligned} E \left[ |Y_t^{(n)} - Y_{t_1}^{(n)}|^2 |Y_{t_2}^{(n)} - Y_t^{(n)}|^2 \right] &\leq E \left[ |Y_t^{(n)} - Y_{t_1}^{(n)}|^4 \right]^{1/2} E \left[ |Y_{t_2}^{(n)} - Y_t^{(n)}|^4 \right]^{1/2} \\ &\leq C|t - t_1| |t_2 - t| \leq C|t_2 - t_1|^2. \end{aligned}$$

For distinct  $i, j, k$  and  $l$ ,  $E[(\xi_i)^3 \xi_j] = E[(\xi_i)^2 \xi_j \xi_k] = E[\xi_i \xi_j \xi_k \xi_l] = 0$ . Hence,  $E[|Y_t^{(n)} - Y_s^{(n)}|^4] = n^{-2} E[\{\sum_{i=1}^{\lfloor nt \rfloor} (y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{i}{n})) \xi_i\}^4]$  is

$$\begin{aligned} &E[(\xi_1)^4] n^{-2} \sum_{i=1}^{\lfloor nt \rfloor} \{y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{i}{n})\}^4 \\ &+ \frac{6}{n^2} E[(\xi_1)^2]^2 \sum_{1 \leq i < j \leq \lfloor nt \rfloor} \{y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{i}{n})\}^2 \{y(\frac{\lfloor nt \rfloor}{n}, \frac{j}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{j}{n})\}^2 \\ &= (I_1 + I_2) E[(\xi_1)^4] + 6(J_1 + J_2 + J_3) E[(\xi_1)^2]^2 \end{aligned}$$

for  $t, s$  and  $n$  satisfying (2.6), where

$$I_1 := n^{-2} \sum_{i=1}^{\lfloor ns \rfloor} \{y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{i}{n})\}^4, \quad I_2 := n^{-2} \sum_{i=\lfloor ns \rfloor + 1}^{\lfloor nt \rfloor} y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n})^4$$

and

$$\begin{aligned} J_1 &:= n^{-2} \sum_{(i,j) \in \Lambda_1} \{y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{i}{n})\}^2 \{y(\frac{\lfloor nt \rfloor}{n}, \frac{j}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{j}{n})\}^2, \\ J_2 &:= n^{-2} \sum_{(i,j) \in \Lambda_2} \{y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - y(\frac{\lfloor ns \rfloor}{n}, \frac{i}{n})\}^2 y(\frac{\lfloor nt \rfloor}{n}, \frac{j}{n})^2, \\ J_3 &:= n^{-2} \sum_{(i,j) \in \Lambda_3} y(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n})^2 y(\frac{\lfloor nt \rfloor}{n}, \frac{j}{n})^2 \end{aligned}$$

with  $\Lambda_1 = \{(i, j) : 1 \leq i < j \leq \lfloor ns \rfloor\}$ ,  $\Lambda_2 = \{(i, j) : 1 \leq i \leq \lfloor ns \rfloor, \lfloor ns \rfloor < j \leq \lfloor nt \rfloor\}$ , and  $\Lambda_3 = \{(i, j) : \lfloor ns \rfloor < i < j \leq \lfloor nt \rfloor\}$ . By (2.6),  $\lfloor nt \rfloor - \lfloor ns \rfloor \leq nt - ns + 1 = n(t - s + \frac{1}{n}) \leq 2n(t - s)$ , so that  $\#\Lambda_1 \leq Cn^2$ ,  $\#\Lambda_2 \leq Cn^2(t - s)$ , and  $\#\Lambda_3 \leq Cn^2(t - s)^2$ . Therefore, using (2.1), we have  $|J_i| \leq C(t - s)^2$  for  $i = 1, 2, 3$  and  $t, s$  and  $n$  satisfying (2.6). Similarly,  $|I_i| \leq C(t - s)^2$  for  $i = 1, 2$ . Thus (2.5) follows.  $\square$

Denote by  $\Delta X$  and  $[X]$  the jump and quadratic variation processes of a process  $X$ , respectively, i.e.,  $\Delta X_t := X_t - \lim_{s \uparrow t} X_s$ ,  $[X]_t := \sum_{s \leq t} (\Delta X_s)^2$ .

**Theorem 2.2.** *The process  $\Delta Y^{(n)}$  converges to zero in probability, while  $[Y^{(n)}]$  converges to the deterministic process  $(t)_{0 \leq t \leq T}$  in probability.*

*Proof.* By (2.5) with (2.6),  $E[(\Delta Y_t^{(n)})^4] \leq E[(Y_t^{(n)} - Y_{t-(1/n)}^{(n)})^4] \leq Cn^{-2}$ , so that,

$$E \left[ \sup_{0 \leq t \leq T} (\Delta Y_t^{(n)})^4 \right] \leq E \left[ \sum_{t \leq T} (\Delta Y_t^{(n)})^4 \right] = \sum_{t \leq T} E \left[ (\Delta Y_t^{(n)})^4 \right] \leq C \frac{nT}{n^2} \rightarrow 0$$

as  $n \rightarrow \infty$ . Thus  $\Delta Y^{(n)}$  converges to zero in probability.

We put  $Z_t^{(n)} := \int_0^t z(\frac{\lfloor nt \rfloor}{n}, s) dW_s^{(n)}$  for  $0 \leq t \leq T$ . Then we have  $Y_t^{(n)} = W_t^{(n)} - Z_t^{(n)}$ , whence  $[Y^{(n)}]_t = [W^{(n)}]_t - 2 \sum_{s \leq t} (\Delta W_s^{(n)}) (\Delta Z_s^{(n)}) + [Z^{(n)}]_t$ .

Since  $z(u, u) = 0$ , we have

$$Z_t^{(n)} - Z_{t-\frac{1}{n}}^{(n)} = \frac{1}{\sqrt{n}} \sum_{i=1}^{\lfloor nt \rfloor - 1} \{z(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - z(\frac{\lfloor nt \rfloor - 1}{n}, \frac{i}{n})\} \xi_i \quad (= 0 \quad \text{if } \lfloor nt \rfloor = 1).$$

From this and (2.1),  $E[(\Delta Z_t^{(n)})^2]$  is at most

$$E \left[ (Z_t^{(n)} - Z_{t-\frac{1}{n}}^{(n)})^2 \right] = \frac{1}{n} \sum_{i=1}^{\lfloor nt \rfloor - 1} \{z(\frac{\lfloor nt \rfloor}{n}, \frac{i}{n}) - z(\frac{\lfloor nt \rfloor - 1}{n}, \frac{i}{n})\}^2 \leq \frac{nT}{n} \cdot \frac{C^2}{n^2} = \frac{C}{n^2}.$$

Since  $[Z^{(n)}]_t$  is increasing, we see that

$$E \left[ \sup_{0 \leq t \leq T} [Z^{(n)}]_t \right] = E \left[ [Z^{(n)}]_T \right] = \sum_{t \leq T} E \left[ (\Delta Z_t^{(n)})^2 \right] \leq nT \frac{C}{n^2} = \frac{C}{n}. \quad (2.7)$$

Thus  $[Z^{(n)}]$  converges to zero in probability.

We have  $[W^{(n)}]_t - t = (\lfloor nt \rfloor / n) - t + (1/n) \sum_{i=1}^{\lfloor nt \rfloor} \{(\xi_i)^2 - 1\}$ . Let  $\epsilon > 0$ . Then, by Kolmogorov's inequality (see, e.g. Williams, 1991, Section 14.6),

$$\begin{aligned} P \left( \sup_{0 \leq t \leq T} \frac{1}{n} \left| \sum_{i=1}^{\lfloor nt \rfloor} \{(\xi_i)^2 - 1\} \right| \geq \epsilon \right) &= P \left( \sup_{0 \leq t \leq T} \left| \sum_{i=1}^{\lfloor nt \rfloor} \{(\xi_i)^2 - 1\} \right| \geq n\epsilon \right) \\ &\leq \frac{1}{\epsilon^2 n^2} \sum_{i=1}^{\lfloor nT \rfloor} E \left[ (\xi_i^2 - 1)^2 \right] \leq \frac{nT}{\epsilon^2 n^2} E \left[ (\xi_1^2 - 1)^2 \right] \rightarrow 0 \end{aligned}$$

as  $n \rightarrow \infty$ . From this and the fact that  $0 \leq t - (\lfloor nt \rfloor / n) < 1/n$ , we see that  $[W^{(n)}]$  converges to the deterministic process  $(t)$  in probability.

By Schwarz's inequality, we have

$$\left| \sum_{s \leq t} (\Delta W_s^{(n)}) (\Delta Z_s^{(n)}) \right| \leq [W^{(n)}]_t^{1/2} [Z^{(n)}]_t^{1/2} \leq [W^{(n)}]_T^{1/2} [Z^{(n)}]_T^{1/2},$$

whence, by (2.7),

$$\begin{aligned} E \left[ \sup_{0 \leq t \leq T} \left| \sum_{s \leq t} (\Delta W_s^{(n)}) (\Delta Z_s^{(n)}) \right| \right] &\leq E \left[ [W^{(n)}]_T^{1/2} [Z^{(n)}]_T^{1/2} \right] \\ &\leq E \left[ [W^{(n)}]_T \right]^{1/2} E \left[ [Z^{(n)}]_T \right]^{1/2} \leq T^{1/2} \cdot (Cn^{-1})^{1/2} = Cn^{-1/2}. \end{aligned}$$

Thus the process  $(\sum_{s \leq t} (\Delta W_s^{(n)}) (\Delta Z_s^{(n)}))$  also converges to zero in probability. Combining, we see that  $[Y^{(n)}]$  converges to  $(t)$  in probability.  $\square$

### 3. APPROXIMATING BINARY MARKET

Let  $Y$  be as defined in Section 2. For  $T, \sigma \in (0, \infty)$  and a deterministic continuous function  $b(\cdot)$  on  $[0, T]$ , we consider the stock price process  $S$  that is governed by the following more general stochastic differential equation than (1.1):

$$dS_t = S_t \{b(t)dt + \sigma dY_t\}, \quad 0 \leq t \leq T,$$

where the initial value  $S_0$  is a positive constant. The solution  $S$  is given by  $S_t = S_0 \exp \left\{ \sigma Y_t + \int_0^t b(s)ds - \frac{1}{2} \sigma^2 t \right\}$ . For  $n = 1, 2, \dots$ , let  $Y^{(n)}$  is as in Section 2. We consider the process  $S^{(n)} = (S_t^{(n)})_{0 \leq t \leq T}$  defined by

$$S_t^{(n)} := \prod_{s \leq t} \left\{ 1 + \sigma \Delta Y_s^{(n)} + \frac{1}{n} b\left(\frac{\lfloor ns \rfloor}{n}\right) \right\}, \quad 0 \leq t \leq T.$$

The aim of this section is to prove that  $S^{(n)}$  converges weakly to the process  $S$ .

As in Eq. (10) and (11) of Sottinen (2001), we put

$$Y_t^{(1,n)} := \sum_{s \leq t} \Delta Y_s^{(n)} \mathbf{1}_{\{|\Delta Y_s^{(n)}| < \frac{1}{2} \sigma^{-1}\}}, \quad Y_t^{(2,n)} := \sum_{s \leq t} \Delta Y_s^{(n)} \mathbf{1}_{\{|\Delta Y_s^{(n)}| \geq \frac{1}{2} \sigma^{-1}\}}.$$

Then we have

$$Y_t^{(n)} = Y_t^{(1,n)} + Y_t^{(2,n)}, \quad (3.1)$$

$$[Y^{(1,n)}]_t = \sum_{s \leq t} (\Delta Y_s^{(n)})^2 \mathbf{1}_{\{|\Delta Y_s^{(n)}| < \frac{1}{2} \sigma^{-1}\}}, \quad (3.2)$$

$$[Y^{(2,n)}]_t = \sum_{s \leq t} (\Delta Y_s^{(n)})^2 \mathbf{1}_{\{|\Delta Y_s^{(n)}| \geq \frac{1}{2} \sigma^{-1}\}}, \quad (3.3)$$

$$[Y^{(n)}]_t = [Y^{(1,n)}]_t + [Y^{(2,n)}]_t. \quad (3.4)$$

**Lemma 3.1.** *The process  $[Y^{(2,n)}]$  converges to zero in probability, whence  $[Y^{(1,n)}]$  converges to the deterministic process  $(t)$  in probability. The process  $Y^{(2,n)}$  converges to zero in probability, whence  $Y^{(1,n)}$  converges weakly to  $Y$ .*

*Proof.* By Theorem 2.2, the process  $\Delta Y^{(n)}$  converges to zero in probability. Since (3.3) implies that  $P(\sup_{0 \leq t \leq T} [Y^{(2,n)}]_t \geq \epsilon)$  is at most  $P(\sup_{0 \leq t \leq T} [Y^{(2,n)}]_t > 0) = P(\sup_{0 \leq t \leq T} |\Delta Y_t^{(n)}| \geq \frac{1}{2} \sigma^{-1})$ ,  $[Y^{(2,n)}]$  converges to zero in probability. Therefore, by Theorem 2.2 and (3.4),  $[Y^{(1,n)}]$  converges to zero in probability.

In the same way, since  $P(\sup_{0 \leq t \leq T} |Y_t^{(2,n)}| \geq \epsilon) \leq P(\sup_{0 \leq t \leq T} |\Delta Y_t^{(n)}| \geq \frac{1}{2} \sigma^{-1})$ , it follows from Theorem 2.2 that  $Y^{(2,n)}$  converges to zero in probability. Therefore, by Theorem 2.1, (3.1) and Theorem 4.1 of Billingsley (1968),  $Y^{(1,n)}$  converges weakly to  $Y$ .  $\square$

**Theorem 3.2.** *The process  $S^{(n)}$  converges weakly to  $S$ .*

*Proof.* Write  $S_t^{(n)} = S_t^{(1,n)} S_t^{(2,n)}$ , where  $S_t^{(2,n)} := \prod_{s \leq t} \{1 + \sigma \Delta Y_s^{(2,n)}\}$ ,  $S_t^{(1,n)} := \prod_{s \leq t} \{1 + \sigma \Delta Y_s^{(1,n)} + (1/n)b(\lfloor ns \rfloor/n)\}$ , and the processes  $Y^{(i,n)}$  are as above. We claim the following: (i)  $S^{(1,n)}$  converges weakly to  $S$ ; (ii)  $S^{(2,n)}$  converges to one in probability.

The claim (ii) implies that  $S^{(1,n)}(S^{(2,n)} - 1)$  converges to zero in probability (see Problem 1 in Billingsley, 1968, p. 28). Since  $S_t^{(n)} = S_t^{(1,n)}(S_t^{(2,n)} - 1) + S_t^{(1,n)}$ , we

see from (i) and Theorem 4.1 of Billingsley (1968) that  $S^{(n)}$  converges weakly to  $S$ , as desired.

For  $\epsilon > 0$ , we have  $P(\sup_{0 \leq t \leq T} |S_t^{(2,n)} - 1| \geq \epsilon) \leq P(\sup_{0 \leq t \leq T} |\Delta Y_t^{(n)}| > \frac{1}{2}\sigma^{-1})$ . Since the process  $\Delta Y^{(n)}$  converges to zero in probability by Theorem 2.2,  $S^{(2,n)}$  converges to one in probability. Thus (ii) follows.

We prove (i). Since the exponential is a continuous functional in the Skorohod topology, it is enough to prove that  $\log S^{(1,n)}$  converges weakly to the process  $(\sigma Y_t + \int_0^t b(s)ds - \frac{1}{2}\sigma^2 t)$ . Notice that  $|\sigma \Delta Y_t^{(1,n)}| + \frac{1}{n}|b(\frac{\lfloor nt \rfloor}{n})| < \frac{3}{4}$  for sufficiently large  $n$  and  $t \in [0, T]$ , whence the logarithm  $\log S^{(1,n)}$  is well defined for such  $n$ .

We have  $\log(1+x) = x - \frac{1}{2}x^2 + r(x)x^3$  for  $|x| < 1$ , where  $r(x)$  is a bounded function on  $|x| \leq \frac{3}{4}$ . Hence

$$\begin{aligned} \log S_t^{(1,n)} &= \sum_{s \leq t} \left\{ \sigma \Delta Y_s^{(1,n)} + \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) - \frac{1}{2} \left( \sigma \Delta Y_s^{(1,n)} + \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) \right)^2 \right. \\ &\quad \left. + r \left( \sigma \Delta Y_s^{(1,n)} + \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) \right) \cdot \left( \sigma \Delta Y_s^{(1,n)} + \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) \right)^3 \right\} \\ &= \sigma Y_t^{(1,n)} + \sum_{s \leq t} \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) - \frac{1}{2}\Phi_t^{(n)} + \Psi_t^{(n)}, \end{aligned}$$

where  $\Phi_t^{(n)} := \sum_{s \leq t} \{ (1/n)b(\lfloor ns \rfloor/n) + \sigma \Delta Y_s^{(1,n)} \}^2$  and

$$\Psi_t^{(n)} := \sum_{s \leq t} r \left( \sigma \Delta Y_s^{(1,n)} + \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) \right) \cdot \left( \sigma \Delta Y_s^{(1,n)} + \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right) \right)^3.$$

We put  $\Gamma_t^{(n)} := \sum_{s \leq t} \frac{1}{n}b\left(\frac{\lfloor ns \rfloor}{n}\right)\Delta Y_s^{(1,n)}$ . Then

$$\Phi_t^{(n)} = n^{-2} \sum_{s \leq t} b\left(\frac{\lfloor ns \rfloor}{n}\right)^2 + 2\sigma \Gamma_t^{(n)} + \sigma^2 [Y^{(1,n)}]_t.$$

Since  $b(\cdot)$  is bounded, the first term  $n^{-2} \sum_{s \leq t} b\left(\frac{\lfloor ns \rfloor}{n}\right)^2$  goes to 0 as  $n \rightarrow \infty$ . By Lemma 3.1, the third term  $\sigma^2 [Y^{(1,n)}]$  converges to  $(\sigma^2 t)$  in probability. As for the second term, it holds that

$$\sup_{0 \leq t \leq T} |\Gamma_t^{(n)}| \leq C \sup_{s \leq T} |\Delta Y_s^{(1,n)}| \leq C \leq |\Delta Y_t^{(n)}|.$$

Since  $\Delta Y^{(n)}$  converges to zero in probability by Theorem 2.2, so does  $\Gamma^{(n)}$ . Thus the process  $(\Phi_t)$  converges to  $(\sigma^2 t)$ . Since  $\sup_{0 \leq t \leq T} \Psi_t \leq C(\frac{1}{n} + \sup_{s \leq T} |\Delta Y_s^{(1,n)}|)\Phi_T$ , we see that the process  $(\Psi_t)$  converges to zero in probability. Using these facts, Lemma 3.1, and Theorem 4.1 of Billingsley (1968), we see that  $\log S^{(1,n)}$  converges weakly to  $(\sigma Y_t + \int_0^t b(s)ds - \frac{1}{2}\sigma^2 t)$ .  $\square$

#### 4. ARBITRAGE OPPORTUNITIES IN THE BINARY MARKET

In this section, we study the arbitrage opportunities in the approximating binary market model with memory constructed in Section 3. For simplicity, we assume that the function  $b(\cdot)$  is a real constant as in (1.1).

Let  $N \in \mathbf{N}$ ,  $r, b \in \mathbf{R}$ , and  $\sigma \in (0, \infty)$ . The number  $N$  corresponds to  $n$  in Sections 2 and 3. Let  $y(t, u)$  be as in Section 2. We define  $r^{(N)} := r/N$ ,  $b^{(N)} := b/N$ . The  $[NT]$ -period market  $\mathcal{M}^{(N)}$  consists of a share of the money market with price



process  $(B_n^{(N)})_{n=0,1,\dots,[NT]}$  and a stock with price process  $(S_n^{(N)})_{n=0,1,\dots,[NT]}$ . The prices are governed respectively by

$$\begin{aligned} B_0^{(N)} &= 1, & B_n^{(N)} &= B_{n-1}^{(N)}(1 + r^{(N)}), & n &= 1, \dots, [NT], \\ S_0^{(N)} &= s_0, & S_n^{(N)} &= S_{n-1}^{(N)}(1 + b^{(N)} + X_n^{(N)}), & n &= 1, \dots, [NT], \end{aligned}$$

where  $s_0$  is a positive constant,

$$X_n^{(N)} := \sigma \Delta Y_{\frac{n}{N}}^{(N)} = \frac{\sigma}{\sqrt{N}} \sum_{i=1}^n \{y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})\} \xi_i$$

and  $\{\xi_i\}$  are i.i.d. random variables such that  $P(\xi_1 = 1) = P(\xi_1 = -1) = 1/2$ . By Theorem 3.2, the binary market model  $\mathcal{M}^{(N)}$  approximates the continuous-time market model with bond price process  $(e^{rt})$  and stock price process  $S$  in (1.1).

Given the values of  $\xi_1, \dots, \xi_{n-1}$ , the random variable  $X_n^{(N)}$  takes the following two possible values  $u_n$  and  $d_n$ :  $d_1 = -\sigma/\sqrt{N}$ ,  $u_1 = \sigma/\sqrt{N}$ , and for  $n = 2, \dots, N$ ,

$$\begin{aligned} d_n &\equiv d_n(\xi_1, \dots, \xi_{n-1}) = \frac{\sigma}{\sqrt{N}} \sum_{i=1}^{n-1} \{y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})\} \xi_i - \frac{\sigma}{\sqrt{N}}, \\ u_n &\equiv u_n(\xi_1, \dots, \xi_{n-1}) = \frac{\sigma}{\sqrt{N}} \sum_{i=1}^{n-1} \{y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})\} \xi_i + \frac{\sigma}{\sqrt{N}}. \end{aligned}$$

We investigate the arbitrage opportunities in  $\mathcal{M}^{(N)}$ . Choose  $C \in (0, \infty)$  so that

$$|y(t, u) - y(s, u)| \leq C|t - s|, \quad 0 \leq t, s, u \leq T. \quad (4.1)$$

**Theorem 4.1.** *Suppose that  $T < 1/C$ . Then there exists an integer  $N_0$  such that for each  $N \geq N_0$ , the market  $\mathcal{M}^{(N)}$  is arbitrage-free.*

*Proof.* From the condition  $TC < 1$ , we have an integer  $N_0$  such that

$$\frac{b}{N} - \frac{\sigma}{\sqrt{N}}(TC + 1) > -1, \quad |r - b| < \sqrt{N}(1 - TC)\sigma \quad (4.2)$$

if  $N \geq N_0$ . Let  $n \in \{1, \dots, [NT]\}$ . Then, by (4.1),  $\min_{\xi \in \{-1, 1\}^{n-1}} d_n(\xi) = -\frac{\sigma}{\sqrt{N}} \sum_{i=1}^{n-1} |y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})| - \frac{\sigma}{\sqrt{N}}$  is at least

$$-\sigma N^{-1/2} [\{(n-1)C/N\} + 1] \geq -\sigma N^{-1/2} (TC + 1).$$

This and (4.2) yield  $b^{(N)} + X_n^{(N)} \geq (b/N) + \min_{\xi \in \{-1, 1\}^{n-1}} d_n(\xi) > -1$  for  $N \geq N_0$  and  $n = 1, \dots, [NT]$ , whence  $S_n > 0$ .

We show that  $\mathcal{M}^{(N)}$  is arbitrage-free for  $N \geq N_0$ . By Proposition 6.1.2 of Dzhaparidze (1996),  $\mathcal{M}^{(N)}$  is free from arbitrage opportunities if and only if

$$d_n < r^{(N)} - b^{(N)} < u_n, \quad n = 1, \dots, [NT]. \quad (4.3)$$

However,  $\max_{\xi \in \{-1, 1\}^{n-1}} d_n(\xi) = \sigma N^{-1/2} \sum_{i=1}^{n-1} |y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})| - \sigma N^{-1/2}$  is at most  $-\sigma N^{-1/2} [1 - \{(n-1)C/N\}] \leq -\sigma N^{-1/2} (1 - TC)$ . Similarly, we see that  $\min_{\xi \in \{-1, 1\}^{n-1}} u_n(\xi) = -\sigma N^{-1/2} \sum_{i=1}^{n-1} |y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})| + \sigma N^{-1/2}$  is at least  $\sigma N^{-1/2} [1 - \{(n-1)C/N\}] \geq \sigma N^{-1/2} (1 - TC)$ . Thus, by (4.2), (4.3) holds for  $N \geq N_0$ .  $\square$

By Theorem 4.1, the market  $\mathcal{M}^{(N)}$  is arbitrage-free for  $T$  small enough and  $N$  large enough. However, in general, the market  $\mathcal{M}^{(N)}$  may admit arbitrage opportunities, as we see below.

Suppose that there exists a positive constant  $C$  such that  $l(s, u) \geq C$  for  $0 \leq u < s \leq T$ . Let  $T > 1/C$ . We assume that  $r \leq b$ . Then,  $d_{\lfloor NT \rfloor}(-1, \dots, -1)$  is

$$\frac{\sigma}{\sqrt{N}} \sum_{i=1}^{\lfloor NT \rfloor - 1} \int_{\frac{i}{N}}^{\frac{i+1}{N}} l(s, \frac{i}{N}) ds - \frac{\sigma}{\sqrt{N}} > \frac{\sigma}{\sqrt{N}} \left( \frac{C(\lfloor NT \rfloor - 1)}{N} - 1 \right).$$

Since  $TC > 1$ , we have  $d_{\lfloor NT \rfloor}(-1, \dots, -1) > r_N - b_N$  or  $S_{\lfloor NT \rfloor} > (1 + r_N)S_{\lfloor NT \rfloor - 1}$  for  $N$  large enough. Therefore, if the value of  $(\xi_1, \dots, \xi_{\lfloor NT \rfloor - 1})$  turns out to be  $(-1, \dots, -1)$ , then we have an arbitrage opportunity: we may buy stocks at time  $\lfloor NT \rfloor - 1$  using money obtained by shortselling bonds. In a similar fashion, we can show that if  $T > 1/C$ ,  $r < b$  and  $N$  is large enough, then the value  $(1, \dots, 1)$  of  $(\xi_1, \dots, \xi_{\lfloor NT \rfloor - 1})$  gives an arbitrage opportunity.

Put  $P_N = P(\bigcup_{n=1}^{\lfloor NT \rfloor} \{d_n < r^{(N)} - b^{(N)} < u_n\}^c)$ . As we see in the proof of Theorem 4.1, the binary market  $\mathcal{M}^{(N)}$  is arbitrage-free if and only if  $P_N = 0$ . The next theorem gives the rate at which the arbitrage probability  $P_N$  tends to zero.

**Theorem 4.2.** *There exists a positive constant  $C' = C'_T$  such that, for each  $\alpha \in (0, 1)$ , we have  $N(\alpha) \in \mathbf{N}$  satisfying  $P_N \leq C' N^{-\alpha}$  for  $N \geq N(\alpha)$ .*

*Proof.* Set  $\beta := (\alpha + 1)/2$ , and choose  $N(\alpha) \in \mathbf{N}$  so large that

$$N^{\beta/2} C \sqrt{T} < \sqrt{N} - |(r - b)/\sigma|, \quad N^{\beta/2} > 4. \quad (4.4)$$

if  $N \geq N(\alpha)$ . Then  $d_1 < r^{(N)} - b^{(N)} < u_1$ . For  $N \geq N(\alpha)$  and  $n = 2, \dots, \lfloor NT \rfloor$ , we put  $\lambda := N^{\beta/2}$ ,  $M_{n-1} := \max_{1 \leq m \leq n-1} |\sum_{i=1}^m \eta_i|$  and

$$s_{n-1} := \left[ N \sum_{i=1}^{n-1} \left\{ y\left(\frac{n}{N}, \frac{i}{N}\right) - y\left(\frac{n-1}{N}, \frac{i}{N}\right) \right\}^2 \right]^{1/2},$$

where  $\eta_i := \sqrt{N} \{y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})\} \xi_i$  for  $i = 1, 2, \dots$ . By (4.1), we have  $s_{n-1} \leq C\sqrt{T}$ . This and (4.4) imply that  $P((r - b)/N \leq d_n)$  is at most

$$P\left(\sigma^{-1}(r - b) + \sqrt{N} \leq M_{n-1}\right) \leq P(M_{n-1} \geq \lambda C \sqrt{T}) \leq P(M_{n-1} \geq \lambda s_{n-1}).$$

Similarly we have  $P(u_n \leq (r - b)/N) \leq P(M_{n-1} \geq \lambda s_{n-1})$ . Since  $\frac{1}{4}\lambda > 1$  and

$$\max_{1 \leq i \leq n-1} |\eta_i| = \max_{1 \leq i \leq n-1} |\sqrt{N} \{y(\frac{n}{N}, \frac{i}{N}) - y(\frac{n-1}{N}, \frac{i}{N})\}| \leq s_{n-1},$$

it follows from Eq. (12.16) of Billingsley (1968, p. 89) that  $P(M_{n-1} \geq \lambda s_{n-1}) \leq C_0 \lambda^{-4}$  for some constant  $C_0 > 0$  independent of  $N$  and  $n$ ; notice that  $\eta_i$  here corresponds to  $\xi_i$  in Eq. (12.16) of Billingsley (1968, p. 89). Hence,  $P_N$  is at most

$$\sum_{n=2}^{\lfloor NT \rfloor} \left\{ P\left(\frac{r - b}{N} \leq d_n\right) + P\left(u_n \leq \frac{r - b}{N}\right) \right\} \leq \frac{2\lfloor NT \rfloor C_0}{N^{2\beta}} \leq \frac{2TC_0}{N^\alpha}.$$

Thus the theorem follows.  $\square$

## REFERENCES

- Anh, V., Inoue, A., 2005. Financial markets with memory I: Dynamic models. *Stochastic Anal. Appl.* 23, 275–300.
- Anh, V., Inoue, A., Kasahara, Y., 2005. Financial markets with memory II: Innovation processes and expected utility maximization. *Stochastic Anal. Appl.* 23, 301–328.
- Billingsley, P., 1968. *Convergence of Probability Measures*. Chapman & Hall, New York.
- Dzhaparidze, K., 1996. *Introduction to Option Pricing in a Securities Market I: Binary Models*. *CWI Quarterly* 9, 319–355.
- Hida, T., Hitsuda, M., 1991. *Gaussian Processes*. American Mathematical Society, Providence.
- Inoue, A., Nakano, Y., Anh, V., 2006. Linear filtering of systems with memory and application to finance. *J. Appl. Math. Stoch. Anal.* 2006, to appear.
- Liptser, R.S., Shiriyayev, A.N., 2001. *Statistics of Random Processes. I. General Theory*, 2nd ed. Springer, New York.
- Sottinen, T., 2001. Fractional Brownian motion, random walks and binary market models. *Finance Stochast.* 5, 343–355.
- Williams, D., 1991. *Probability with Martingales*. Cambridge University Press, Cambridge.
- E-mail address:* inoue@math.sci.hokudai.ac.jp
- E-mail address:* y-nakano@sigmath.es.osaka-u.ac.jp
- E-mail address:* v.anh@fsc.qut.edu.au

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN

DIVISION OF MATHEMATICAL SCIENCES FOR SOCIAL SYSTEMS, GRADUATE SCHOOL OF ENGINEERING SCIENCE, OSAKA UNIVERSITY, TOYONAKA 560-8531, JAPAN

SCHOOL OF MATHEMATICAL SCIENCES, QUEENSLAND UNIVERSITY OF TECHNOLOGY, GPO Box 2434, BRISBANE, QUEENSLAND 4001, AUSTRALIA